

**2019 Young Investigator Award 2<sup>nd</sup> Runner-Up**

**616: RNA-SEQ ANALYSIS OF BLASTOCOEL FLUID REVEALS A UNIQUE GENE EXPRESSION SIGNATURE PROFILE IN EUPLOID EMBRYOS THAT SUCCESSFULLY IMPLANT**

**Alyssa Barre<sup>1</sup>, Allison Kranyak<sup>1</sup>, Deepti Athavale<sup>1</sup>, Lauren Tyler<sup>1</sup>, Arnav Lal<sup>1</sup>, Shawn Zimmerman<sup>2</sup>, T. Arthur Chang<sup>3</sup>, Randal Robinson<sup>3</sup>, J. David Wininger<sup>4</sup>, William Roudebush<sup>1</sup>, Renee Chosed<sup>1</sup>**

<sup>1</sup> Biomedical Sciences, University of South Carolina School of Medicine Greenville, Greenville, United States of America

<sup>2</sup> VIOS Fertility Institute, VIOS Fertility Institute, Swansea, United States of America

<sup>3</sup> Obstetrics and Gynecology, University of Texas Health Science Center, San Antonio, United States of America

<sup>4</sup> Atlantic Reproductive Medicine Specialists, Atlantic Reproductive Medicine Specialists, Raleigh, United States of America

**Objective**

Discovering a molecular signature in day-5 blastocysts that is suggestive for a successful uterine implantation would provide reproductive specialists an additional tool for selecting the very best embryo for transfer. This study assessed global gene expression using RNA-Seq in blastocoel fluid-conditioned media from euploid embryos resulting in (un)successful implantations.

**Design**

Retrospective analysis of day-5 euploid blastocoel fluid gene expression and implantation outcomes.

**Materials and Methods**

Blastocoel fluid-conditioned media was obtained following biopsy of ICSI-generated day-5 blastocysts. RNA was extracted and libraries prepared using a SMART-Seq Stranded kit followed by Illumina NextSeq500 sequencing. Sequences were aligned to the human genome, reads counted and gene expression determined. The PANTHER classification system (pantherdb.org) was used to identify signaling pathways most represented in the RNA-Seq gene lists per sample. Embryo implantation-related genes were included in the analysis (Sanchez-Ribas et al., Fert Steril 2019;111:991).

**Results**

A greater number of expressed genes (n=1484) were found associated with no euploid implants than embryos (n=778) that did implant. A greater percentage of genes belonging to apoptotic (1.2 vs 0.6%), GnRH (2.4 vs 1.4%), inflammation (3.1 vs 0.8%) and Wnt (2.3 vs 1.5%) signaling pathways were found to be associated with a successful vs unsuccessful implants. These pathways are elevated in the embryo implantation-related genes. The ubiquitin-proteasome signaling pathway had a greater expression percentage in the negative (0.9%) pregnancy outcomes than positive (0.3%) outcomes.

**Conclusions**

We identified specific gene expression in unique signaling pathways in conditioned media from euploid embryos capable of establishing a successful pregnancy outcome.

**Support**

UTHSCSA

**Disclosure**

None